



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

converges only when $x \equiv 1$. Moreover, the given series is not convergent, since

$$\lim_{n \rightarrow \infty} \frac{(e-1)^n}{n} \neq 0,$$

$e - 1$ being greater than unity.

Also solved by HORACE OLSON, C. E. FLANAGAN, J. W. CLAWSON, and H. S. UHLER.

GEOMETRY.

449. Proposed by H. E. TREFETHEN, Colby College.

Find a tetrahedron with the face angles at one vertex in arithmetical progression and its six edges expressed in positive integers.

SOLUTION BY THE PROPOSER.

Let the angles be $A + B$, A , $A - B$; the lateral edges x, y, z ; the base edges a, b, c ; so that $a^2 = x^2 + y^2 - 2xy \cos(A + B)$, $b^2 = x^2 + z^2 - 2xz \cos A$, $c^2 = y^2 + z^2 - 2yz \cos(A - B)$. If, for brevity, we put $1 + \cos(A + B) = P/2$, $1 + \cos A = Q/2$, and $1 + \cos(A - B) = R/2$, we may write

$$a^2 = (x + y)^2 - Pxy = (x + y - p)^2 \text{ or } Pxy - 2py = 2px - p^2, \quad (1)$$

$$b^2 = (x + z)^2 - Qxz = (x + z - q)^2 \quad Qxz - 2qz = 2qx - q^2, \quad (2)$$

$$c^2 = (y + z)^2 - Ryz = (y + z - r)^2 \quad Ryz - 2ry = 2rz - r^2. \quad (3)$$

Eliminating y and z from (1), (2), (3), and arranging we have

$$(2px - p^2)[q(4r - Rq) + 2(Rq - Qr)x] = r(Px - 2p)[2q(r - q) + (4q - Qr)x]. \quad (4)$$

If we put the coefficient of $x^2 = 0$, then

$$p = \frac{Pr(4q - Qr)}{4(Rq - Qr)} \quad (5)$$

and also

$$2x = \frac{p^2q(4r - Rq) + 4pqr(q - r)}{p(8qr - Qr^2 - Rq^2) - p^2(Rq - Qr) + Pqr(q - r)}. \quad (6)$$

We may now assign values to A and B and thus determine P, Q, R . If values are assigned to q and r also, p is defined by (5), and then x may be found from (6), y and z from (1), (2), (3) and finally a, b, c also.

Thus if $A = 90^\circ$ and $B = \arcsin 1/3$, then $P = 4/3$, $Q = 2$, $R = 8/3$; and if also $q = r = 1$, then $p = 1$, $x = 1/4$, $y = 3/10$, $z = 1/3$. Reducing the values of x, y, z to a common denominator and rejecting it, we have in integers $x = 15$, $y = 18$, $z = 20$, and consequently $a = 27$, $b = 25$, $c = 22$. Or since the equations are symmetrical we may use the reciprocals $x = 4$, $y = 10/3$, $z = 3$, whence in integers $x = 12$, $y = 10$, $z = 9$, and then $a = 18$, $b = 15$, $c = 11$. Again if $\sin B = 1/2$, the angles are 120° , 90° , and 60° ; $x = 9$, $y = 15$, $z = 40$; $a = 21$, $b = 41$, $c = 35$. If $\sin B = 2/3$, we find $x = 1,092$, $y = 416$, $z = 81$; $a = 2,804$, $b = 2,185$, $c = 367$.

By varying q, r and the $\sin B$, other sets of numbers may be found ad libitum.

464. Proposed by FRANK R. MORRIS, Glendale, Calif.

The sum of the hypotenuse and one side of a right triangle is 100 feet. A point on the hypotenuse is 10 feet from each of the sides. Find the length of the hypotenuse correct to the third decimal place.

SOLUTION BY W. W. BURTON, Mercer University, Macon, Ga.

Let $BC = x$. Then $AB = 100 - x$ and $BD = x - 10$. The right triangles ACB and PDB are similar. (Their sides are respectively parallel.) Therefore

$$AB : BC = PB : BD \text{ or } (100 - x) : x = PB : (x - 10),$$